

# SUMMER-FALL FESCUE MINERAL SUPPLEMENT

Mineral supplement for beef cattle on fescue pasture from June - Dec **MEDICATED** 

Control of bacterial pneumonia associated with shipping fever complex caused by Pasteurella spp. susceptible to chlortetracycline.

#### **ACTIVE DRUG INGREDIENT**

Chlortetracycline		5000 g/ton (2500 mg/lb)
-	GUARANTEED ANALYSIS	,
Calcium (Ca),	Min.	10.50%
Calcium (Ca),	Max.	12.60%
Phosphorus (P),	Min.	8.00%
Salt (NaCl),	Min.	17.00%
Salt (NaCl),	Max.	20.40%
Magnesium (Mg),	Min.	3.00%
Potassium (K),	Min.	0.15%
Copper (Cu),	Max.	300 ppm
Selenium (Se),	Min.	13 ppm
Zinc (Zn),	Min.	40 ppm
Vitamin A,	Min.	50,000 IU/lb

## **INGREDIENTS**

Dicalcium Phosphate, Mono-Calcium Phosphate, Salt, Magnesium Oxide, Cane Molasses, Vitamin A Supplement, Vitamin D3 Supplement, Vitamin E Supplement, Sulfur, Bentonite, Potassium Sulfate, Magnesium Sulfate, Maganous Oxide, Zinc Oxide, Iron Sulfate, Copper Oxide, Cobalt Carbonate, Ethylene Diamine Dihydriodide, Dried Distillers Grains with Solubles, Dried Lactobacillus Acidophilus Fermentation Product, and Sodium

## FEEDING DIRECTIONS

All other minerals should be removed. Feed S-F Fescue Mineral from June through December at the following rates: feed 2.25 oz. per head per day to beef cattle to provide 350 mg/head/day.

## CAUTION

This product contains added copper sulfate. Do not p noice to sheep that are subject to copper toxicity.

WARNING
A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal. For feeds providing 350 mg chlortetracycline or more per head per day, discontinue use 48 hours before animals are slaughtered. Do not feed to lactating dairy cows.

#### Manufactured by





**Buchheit Premium Feeds** 33 PCR 540 Perryville, MO 63775 1-800-622-7937 (573) 547-4569

NET WEIGHT 50# (22.7 Kg)